FORM PTO-1390 U.S. DEPARTMENT OF COM	MERCE PATENT AND TRADEMARK OFFICE	ATTORNEY 'S DOCKET NUMBER	
(REV. 1-98)  TRANSMITTAL LETTER TO THE UNITED STATES		MSKP039	
TRANSMITTAL LETTER DEGIGNATED/ELECT	ED OFFICE (DO/EO/US)	U.S. APPLICATION NO (If known, see 37 CFR 15	
CONCERNING A FILIN	IG UNDER 35 U.S.C. 371	09/674975	
INTERNATIONAL APPLICATION NO PCT/US99/10065	INTERNATIONAL FILING DATE 07 May 1999	PRIORITY DATE CLAIMED  08 May 1998	
TITLE OF INVENTION COMPOSITIONS AND METHODS FOR	ACTIVE VACCINATION	ı EL556132244U	
APPLICANT(S) FOR DO/EO/US		C	
Agus, et al.  Applicant herewith submits to the United St	ates Designated/Elected Office (DO/EO/US	s) the following items and other information:	
	is concerning a filing under 35 US C 371		
<del></del>	NT submission of items concerning a filing	g under 35 US.C 371	
This express request to begin nation		) at any time rather than delay	
		h month from the earliest claimed priority date	
Little -			
a. is transmitted herewith (re	equired only if not transmitted by the Internation	ational Bureau).	
b. has been transmitted by the	le International Bureau. lication was filed in the United States Recei	iving Office (RO/US)	
c. is not required, as the appl	pplication into English (35 U.S C 371(c)(2)		
	ternational Aplication under PCT Article 19		
Amendments to the claims of the Infanta a. are transmitted herewith (b. have been transmitted by c. have not been made; how d. have not been made and v	required only if not transmitted by the Inter-	national Bureau).	
b. have been transmitted by	the International Bureau		
c. have not been made; however, the time limit for making such amendments has NOT expired			
d. A have not been made and v		251 ( )(2))	
8. A translation of the amendments to the claims under PCT Article 19 (35 U S C 371 (c)(3)).			
9. An oath or declaration of the invent		1 DOM 4 / 1 2 /	
10. A translation of the annexes of the I (35 U.S C 371(c)(5))	International Preliminary Examination Repo	ort under PCT Article 36	
Items 11. to 16 below concern docume	ent(s) or information included:		
11.  An Information Disclosure Stateme			
12. An assignment document for record	ling A separate cover sheet in compliance	with 37 CFR 3 28 and 3.31 is included	
13. A FIRST preliminary amendment	<del>_</del>		
☐ A SECOND or SUBSEQUENT pre	eliminary amendment.		
14. A substitute specification.			
15. A change of power of attorney and	or address letter		
16. Other items or information:		E	
		Exp. Mail No.: EL556/32244US  Date Mailed: OT NOVEMBER 2000  Name: SUSAN HOWARD  Sign:	

## **528** Rec'd PCT/PTO **0** 7 NOV 2000

U.S. APPLICATION NO (15kno)	vi. 5237 CFR (5) 7 K PCT/U	TERNATIONAL APPLICATION NO S99/10065			ATTORNEY'S DOCKE MSKP039	T NUMBER
17. The following fees are submitted:			CAI	LCULATIONS P	TO USE ONLY	
BASIC NATIONAL FEE (37 CFR 1 492 (a) (1) - (5)):				***		
Neither international preliminary examination fee (37 CFR 1.482)						
nor international se and International S	arch fee (37 ČFR 1.445(a earch Report not prepare	a)(2)) paid to USPTO d by the EPO or JPO	\$1070 00			
International prelim USPTO but Interna	ninary examination fee (1 ntional Search Report pre	37 CFR 1 482) not paid to pared by the EPO or JPC	o ) \$930 00			
International prelimbut international se	ninary examination fee (3 arch fee (37 CFR 1 445(	37 CFR 1 482) not paid to a)(2)) paid to USPTO	O USPTO \$790 00			
but all claims did r	not satisfy provisions of I	37 CFR 1.482) paid to US PCT Article 33(1)-(4)	(\$720.00			
International prelin	ninary examination fee (1 fied provisions of PCT A	37 CFR 1.482) paid to U	SPTO \$98.00			
		BASIC FEE AMO		\$	710 00	
			□ 20 □ 30	-	710,00	
Surcharge of \$130.0 months from the ear	0 for furnishing the oath liest claimed priority dat	e (37 CFR 1 492(e))		\$		
CLAIMS	NUMBER FILED	NUMBER EXTRA	RATE	\$		
Total claims	<u> </u>	0	x \$22.00	\$		
Independent claims	4 -3 =	-1:1-1-)	x \$82.00	\$	89.00	
+4427	DENT CLAIM(S) (if app	OF ABOVE CALCU	+ \$270.00 ILATIONS =	\$	79200	
Baduction of 1/2 for		fapplicable. A Small Ent			792,00	
must also be filed (I	Note 37 CFR 1.9, 1.27, 1.	.28) Applicant clair	ms small entity status +	\$ _	396.00	
egia			JBTOTAL =	\$	396,00	
Processing fee of \$1	30.00 for furnishing the	English translation later	than 20 30	\$		
months from the ear	liest claimed priority dat					
TOTAL NATIONAL FEE =			\$	396.00		
Fee for recording the accompanied by an	e enclosed assignment (3 appropriate cover sheet (	7 CFR 1.21(h)) The ass 37 CFR 3 28, 3.31) \$40	.00 per property +	\$		
		TOTAL FEES E	NCLOSED =	\$	396.00	
			Am	ount to be refunded:	\$	
					charged:	\$
a. A check in the amount of \$ 396. to cover the above fees is enclosed.  b. Please charge my Deposit Account No in the amount of \$ to cover the above fees.						
A duplicat	e copy of this sheet is en	closed.				
c. The Commissioner is hereby authorized to charge any additional fees which may be required, or credit any overpayment to Deposit Account No. 15-0.610 A duplicate copy of this sheet is enclosed.						
NOTE: Where an 1.137 (a) or (b)) m	n appropriate time limi nust be filed and granted	t under 37 CFR 1.494 or d to restore the applicat	1.495 has not been ion to pending status	met, s.	a p <b>e</b> ition to reviv	re (37 CFR
SEND ALL CORRESP	ONDENCE TO		min	11 -	u S X	and
I TELEVISIONE VITE			SIGNATI	URE		
	I 1100A 1101A 1100A 110A 140A		Marina	T. L	arson	
02	1121		NAME			
BATENT TO	DEHARK OFFICE		32,038	3		
THE STATE OF			REGISTE	RATIO	N NUMBER	
					-	

25

5

### PCT/US99/10065

### COMPOSITIONS AND METHODS FOR ACTIVE VACCINATION

#### **BACKGROUND OF THE INVENTION**

This application relates to an active vaccine approach to the treatment of cancer and other diseases. The approach is applicable to a number of cancers and diseases, although a preferred embodiment provides an active vaccine for treatment of B cell Non-Hodgkin's Lymphoma (NHL).

NHL is characterized by a clonal proliferation of malignant B cells. The treatment of NHL across a broad spectrum of patients remains a challenge, although numerous therapeutic approaches have been proposed and tried.

The most common therapeutic approach being used today is chemotherapy.

While chemotherapy is effective for some period of time in most patients, a significant percentage of patients are not cured and experience a relapse.

Treatments have been proposed based on anti-idiotype therapy. In anti-idiotype therapy, a cell surface molecule which is expressed by malignant cells but not by normal cells is used to create patient-specific antibodies which are then administered to the patient. See, Miller, et al., *New Engl. J. Med.* 306: 517-522 (1982). Autologous patient-derived idiotype proteins have also been conjugated with keyhole limpet hemocyanin to produce a vaccine which has demonstrated efficacy and can elicit B and T cell immune responses. Kwak et al., *New Engl. J. Med.* 327: 1209-1215 (1992). Hybridoma-derived idiotype was co-cultured with patient-derived dendritic cells which acted as antigen presenters upon re-infusion into the patient and showed clinical efficacy. Hsu et al., *Nature Medicine* 2: 52-58 (1996). Idiotypic vaccines made in lipid-based carriers are disclosed in International Patent Publication WO98/14170.

Treatments have also been proposed using antibodies directed to CD20, a transmembrane protein that is expressed by both normal and malignant B-cells during parts of the B cell development cycle. Using single-dose infusions with anti-CD20 monoclonal antibodies, partial or minor tumor regressions were observed in 6 of 15 patients in a Phase I clinical study. Maloney et al., *Blood* 84: 2457-2466 (1994). In Phase II studies, 17 of 37 patients showed complete or partial remissions. In December 1997, the FDA approved the

first antibody-based therapy for NHL. Rituximab (Ritvaxan, IDEC/Genentech) is a chimeric human/murine antibody approved for the treatment of patients with relapsed or refractory low-grade or follicular CD20<sup>+</sup> B cell NHL. Maloney et al., *Blood* 90: 2188-2195 (1997).

Combinations of chemotherapy and anti-CD20 therapy have been reported as having better therapeutic efficacy, with 11 of 11 patients showing complete or partial remission. Czuczman et al., Abstract 53, *Ann. Oncol.* 7, Supp. 1: 56 (1996).

While therapeutic regimens using anti-CD20 concepts are potentially effective, all of these therapies have the drawback of being passive therapies, i.e., they do not directly involve the immune system of the patient. Thus, these therapies may require the continued administration of the therapeutic agent for efficacy and do not provide any long-term protection against recurrence. In addition, the passive therapy is monoclonal in nature, therefore escape is possible. It would therefore be desirable to have an active therapy, that is a therapeutic agent which when administered to the patient stimulates an immune response against CD20 found in B-cells.

It is an object of the present invention to provide such a therapy. It is a further object of the invention to provide an active polyclonal therapy that is difficult to evade.

#### SUMMARY OF THE INVENTION

20

In accordance with the present invention, NHL is treated, not by administration of an anti-CD20 monoclonal antibody, but by the administration of CD20 itself, or an immunogenic fragment of the extracellular portion thereof, coupled to or administered with an antigenic carrier moiety such as keyhole limpet hemocyanin (KLH). This results in the stimulation of the production of polyclonal antibodies against CD20 (or an immunogenic fragment thereof) which has the affect of reducing the number of B-cells, including malignant B-cells. Thus, the invention provides an active vaccine. The same approach can be used for therapeutics for other diseases and conditions in which target cells possess a transmembrane protein, and is particularly applicable to those diseases and conditions for which administration of antibodies to transmembrane proteins or peptides (i.e., passive therapy) have been shown to provide therapeutic benefits, and especially in the

25

30

5

situations where the target is also capable of transducing or receiving a signal important for cell growth or function. This would include, for example, Her2/neu, VEGF receptor, epidermal growth factor receptor, the CD19 molecule, interleukin-2-receptor, interleukin-4-receptor, and the P-glycoprotein, also known as the multidrug-resistance protein.

BRIEF DESCRIPTION OF THE FIGURES

Figs. 1A and B show ELISA results for formation of antibodies to human and mouse CD20 in vaccinated mice;

Figs. 2A and B shows results for binding of control B1 antibodies or antibodies in plasma from a mouse treated with human CD20-KLH conjugate with Raji B NHL cells;

Fig. 3 shows CP19<sup>+</sup>B cell levels in mice treated with human or mouse CD20-KLH conjugate;

Fig. 4 shows the domain structure of human Her2;

Fig. 5 shows the domain structure of human EGFR;

Figs. 6A-D shows the cross-reactivity of antibodies generated in response to human or mouse CD20 fragments;

Figs. 7A-D show the importance of carrier protein and adjuvant in generating an immune response;

Figs. 8A-D shows the immune response generated using different adjuvants; and

Figs. 9A-I shows CP19<sup>+</sup>B cell levels in mice treated with human or mouse CD20-KLH conjugate.

#### DETAILED DESCRIPTION OF THE INVENTION

The present invention provides an active vaccine therapy which can be used in the treatment of a variety of cancers and related conditions in which it is desirable to bring about the death of a target group of cells. Conventionally, immunotherapies targeting cells have sought to obtain a cellular immune response (T-cells that recognize the target cells), since a humoral immune response (antibodies that recognize the target cells) alone is not

25

30

5

deemed sufficient to achieve the desired result of cell death. The present invention departs from this conventional wisdom, and effectively utilizes a humoral immune response against the target cells to provide therapeutic benefit. The targets for therapy include cell surface proteins that when bound by a ligand signal to the cell. The vaccine induced antibody response will mimic ligand binding and cause similar signaling events which can imitate the process of programmed cell death (apoptosis) or halt the cell from growing or change the cancer cell's sensitivity to chemotherapy.

By way of example, the invention is suitably employed in the treatment of NHL and other B cell diseases such as chronic lymphocytic leukemia, auto-immune disorders and B-cell regulatory disorders. In accordance with this embodiment of the invention, a peptide antigen is prepared which contains at least an immunogenic portion of the extracellular domain of CD20 coupled to or administered with an antigenic carrier protein. The CD20 component of the peptide antigen may be syngeneic or it may be xenogeneic. Thus, for example, human patients may be treated with a peptide vaccine containing a human or a mouse CD20-fragment. There is evidence that strong immune responses can be elicited against xenogeneic proteins. Naftzger et al., Proc. Natl. Acad. Sci. (USA) 93: 14809-14814 (1996): International Patent Application PCT/US97/22669, filed December 10 1997, incorporated herein by reference. A suitable fragment is the 44 amino acid peptide spanning amino acids 136 to 179 of the sequence of mouse or human CD20. (Seq. ID Nos. 1 and 2) Other immunogenic fragments derived from the extracellular domain of CD20, or the entire CD20 molecule may also be used. Seq. ID. Nos. 3 and 4 shows the nucleic acid and amino acid sequences, respectively, of exon VI (the extracellular domain) of human CD20 as reported by Tedder et al., J. Immunol. 142: 2560-2568 (1989).

As used in the specification and claims hereof, an "immunogenic fragment" is a molecule which includes at least a portion of the extracellular domain of a transmembrane protein to direct and immunological response to that transmembrane protein when the immunogenic fragment is coupled to or administered with an antigenic carrier protein effective to break tolerance and administered with an adjuvant. It is not required that the immunogenic fragment alone be effective to stimulate an immune response, although such stimulation would not take a given fragment outside the scope of the present invention.

20

25

A preferred antigenic carrier protein is keyhole limpet hemocyanin which can be coupled to peptides using techniques described in Pierce Catalog Protocol. Other antigenic carrier proteins which can be used to break tolerance might be used in the invention include immunoglobulins, tuberculin, tetanus toxin and others well known in the art.

The peptide antigen containing the CD20 component and the antigenic carrier protein is formulated with a pharmaceutically acceptable adjuvant in a liquid carrier and administered to a patient suffering from NHL or another B cell disease. The composition will generally be administered by injection, for example, intramuscular, subcutaneous or intradermal injection, but might also be administered by way of a DNA vaccine (See US Patent No. 5,580,859, incorporated herein by reference) or a viral vaccine, or after mixing with antigen presenting cells (APC's) such as dendritic cells, ex vivo. Alternatively, the antigen may be administered without adjuvant by injection into a host prepared by prior or simultaneous injection of an immune adjuvant. Specific amounts to be administered to a patient can be determined by monitoring the titer of anti-CD20 antibodies developed by the patient, or by an average group of patients using well-known technology.

When a peptide of the extracellular domain of human or mouse CD20 is coupled to KLH and administered with an adjuvant to mice, antibodies which react with CD20 are found in plasma. (Figs. 1A and B) These antibodies bind to Raji cells, a human lymphoma cell line, indicating the ability to bind to a cell expressing CD20. (Figs. 2A and B). Moreover, the number of CD19<sup>+</sup> B cells present in mice injected with either of the two CD20-KLH conjugates declines substantially (~30% decrease relative to controls). (Figs. 3 and 9). The assay used to quantitate B cell depletion detects CD19 which is also expressed on immature B cells that are CD20<sup>-</sup>. Thus, the 30% depletion actually underestimates the efficacy of the vaccine against CD20<sup>+</sup> B cells. .

Antibodies generated in mice after vaccination with human or mouse-derived CD20 fragments are specific for the peptides used, yet are capable of inducing immunity to the corresponding peptide from other species (Figs. 6A-D). Studies showed that in most instances the peptide, carrier protein and adjuvant are all needed for optimal response, although some responses were detected using less than all of the components. (Figs. 7A-D).

25

30

5

Several different adjuvants were also tested, and QS21 was found to be the most effective of those tested. (Fig. 8A-D).

While not intending to be bound by any particular mechanism, it is believed that the vaccines of the present invention are effective via at least two pathways. First, the generation of a humoral immune response to CD20 is effective to some extent to reduce the numbers of B cells bearing CD20 antigen in a manner consistent with normal immunological response to a target antigen. In addition, however, because CD20 has a signaling function, the binding of antibody to the CD20 moiety activates this signaling function to trigger apoptotic cell death. Such stimulation of apoptosis has been demonstrated to occur *in vitro* following passive treatments with a chimeric anti-CD20 antibody. Maloney et al., *Blood* 88 (Supp. 1): 637a (1996).

It is also possible that T cell mediated effector mechanisms are involved in the immune response. As evidence of this, we illustrate in Table 1 the mouse and human peptide sequences capable of binding to the corresponding mouse and human histocompatability antigens. This information was derived from a search of the NIH Bioinformatics and Molecular Analysis Section HLA Binding Predictions database using the mouse and human CD20 amino acid sequences. (Parker et al., *J. Immunol.* 152: 163 (1994)).

While the method of the invention is illustrated here using CD20 or CD20-derived peptides as the antigen to target B cells, the invention is not limited to this embodiment. Rather, the inventions encompasses the use of vaccine compositions comprising an immunogenic portions of the extracellular domain of transmembrane protein or peptide, particularly a transmembrane protein or peptide having signaling function, coupled to or administered with an antigenic protein and/or adjuvant to break tolerance.

A non-limiting example of another transmembrane protein which can be used in whole or in part in the method of the invention is Her-2/neu. The Her-2/neu oncogene is a receptor-like tyrosine kinase that is expressed on the cell surface of a significant portion of solid tumors. It has been shown that patients with early stage breast cancer have a high titer of antibodies to Her-2/neu. Disis et al., *J. Clin. Oncol.* 15: 3363-3367 (19967). The amino acid sequence and domain structure of human Her-2/neu are shown in Seq. OD. No. 5 and Fig. 4, and isolation and expression of the extracellular domain has been disclosed.

25

5

International Patent Publication No. WO 90/14357, which is incorporated herein by reference. There is clinical data showing efficacy of monoclonal antibodies against Her-2-neu in the treatment of patients with Her-2/neu<sup>+</sup> tumors, and potential synergism with chemotherapy. Thus, in accordance with the present invention, a vaccine composition comprising at least an immunogenic portion of the extracellular domain of Her-2-neu (amino acids 22 to 652) coupled to or administered with an antigenic protein or peptide such a KLH can be used as a vaccine to provide the same therapeutic benefits using an active as opposed to a passive approach.

A further non-limiting example of a transmembrane protein which can be used in whole or in part in the method of the invention is epidermal growth factor receptor (EGFR). The amino acid sequence and domain structure of human EGFR are shown in Seq. ID. No. 6 and Fig. 5. There is significant data showing that antibodies to EGFR can have anti-tumor activity in breast and prostate cancer, as well as several head and neck tumors. Prewett et al., *J. Immunother. Emphasis Tumor Humoral* 19: 419-27 (1996). The mechanism by which antibody therapy against EGFR may be efficacious can be through the ability to down-regulate vascular endothelial growth factor production by tumor cells and thereby decrease angiogenesis. Petit et al., *Am. J. Pathol.* 151: 1523-30 (1997). In accordance with the present invention, a vaccine composition comprising at least an immunogenic portion of the extracellular domain of EGFR (amino acids 25 to 645) coupled to or administered with an antigenic protein or peptide such a KLH can be used as a vaccine to provide the same therapeutic benefits using an active as opposed to a passive approach. Preferred immunogenic peptides would be selected from regions not deleted in the various types of truncated EGFR mutants associated with some cancers.

A further non-limiting example of a transmembrane protein which can be used in whole or in part in the method of the invention is VEGF receptor. There are significant data showing that antibodies to VEGF receptor can inhibit angiogenesis and thereby halt tumor progression. In accordance with the present invention, a vaccine composition comprising at least an immunogenic portion of the extracellular domain of VEGF receptor coupled to or administered with an antigenic protein or peptide such a KLH can be used as a

vaccine to provide the same therapeutic benefits using an active as opposed to a passive approach.

Still a further non-limiting example of a transmembrane protein which can be used in whole or in part in the method of the invention is the IL-2 receptor. The IL-2 receptor is expressed on most T-cells malignancies, and there is a data showing that antibodies to the IL-2 receptor can be used in the treatment of T-cell malignancies and autoimmune disorders. In the present invention, a composition is made comprising at least an immunogenic portion of the extracellular domain of the IL-2 receptor (e.g., P55 or P75), coupled to or administered with an antigenic carrier protein or peptide such as KLH. and used as a vaccine.

The vaccine compositions of invention can be used alone or in combination (concurrently or sequentially) with drugs or chemotherapy agents that provide therapeutic benefit for the condition being treated. In the case of NHL, suitable chemotherapy agents which can be used in combination with the CD20 based vaccine include alkylating agents, anthrocyclines, cis-platinum, fludarabine, corticosteroids and vinca alkaloids. These same chemotherapy agents which might be used in combination with other vaccine compositions for other forms of cancer.

#### EXAMPLE 1

20

44 amino acid fragments of the extracellular domains of humans and murine CD20 (amino acids 136-179, Seq. ID Nos. 1 and 2) were synthesized using a solid-phase FMOC peptide synthesizer and coupled to KLH using the methodology described in the Pierce Catalog Protocol. The peptide coupled to KLH was then prepared for injection by formulation with QS-21 adjuvant. Balb/c mice were injected according to one of the following protocols on days 1, 8, 15, 22 and 50 of the experiment:

- A. Murine CD20 fragment-KLH with QS-21 adjuvant
- B. Human CD20 fragment-KLH with QS-21 adjuvant
- C. KLH with QS-21 adjuvant
- D. QS-21 adjuvant

5

- E. P190 (irrelevant protein) coupled to KLH with QS-21 adjuvant
- F. B3A2 (irrelevant peptide) coupled to KLH with QS-21 adjuvant.

The animals were sacrificed on day 62 of the experiment.

Serum samples from the mice were diluted 1:200 and evaluated by BSA-blocked ELISA using goat-anti-mouse antibody conjugated to alkaline phosphatase for antibodies which bind to human CD20, mouse CD20 and KLH. As shown in Figs 1A and B, mice injected with human CD20 coupled to KLH (Fig. 1A) or mouse CD20 coupled to KLH (Fig. 1B) administration of xenogeneic antibody produced a significant polyclonal antibody response to both human and mouse CD20, while the response following administration of syngeneic antibody was principally limited to antibodies to the syngeneic form of CD20. Either xenogeneic or syngeneic peptide can therefore be used to generate an immune response.

To confirm the ability of the antibodies to bind to B cells, Raji cells (a form of human B-cell lymphoma that expresses CD20 on its surface) were blocked with human IgG, washed and then incubated for 30 minutes on ice with a 1:10 dilution of plasma from a mouse vaccinated with P-190-KLH control or huCD20-KLH. As a positive control, Raji cells were incubated with B1 antibody, or IgG2 as an isotypic negative control. After washing, the cells were incubated with goat-anti-mouse antibody, washed and fixed with 1% paraformaldehyde. Flow cytometry analysis was performed in a Becton-Dickinson FACScan. The results are shown in Figs 2A and 3B, wherein the shaded data set are the experimental data set and the outlined data set is the negative controls. As is apparent, there is a strong binding of mouse antibodies and Raji cells, comparable to that observed with B1 antibody.

25 <u>EXAMPLE 2</u>

To assess the number of B cells present in vaccinated mice, an evaluation was made of cells expressing CD19, a standard phenotypic marker for B cells. Spleens were harvested from the animals vaccinated in Example 1 and put into a single-cell suspension.

After counting the total number of cells, the cells were stained with FITC-labeled anti-mouse CD19 and the samples were analyzed by flow cytometry with a FACScan. 10,000 events

were collected. The percentage of CD19 positive cells minus the control gate was multiplied by the total number of cells to determine the number of CD19 positive cells in mice treated with the mouse and human CD20 peptide conjugates, and the P190 irrelevant peptide conjugate control.

5

As shown in Fig. 3, the absolute number of CD19 positive cells was significantly reduced in mice treated with either of the CD20 peptide conjugates. The level of CD19 positive cells is a reflection of the number of CD20 positive B cells, and the number of immature CD19<sup>+</sup>, CD20<sup>-</sup> B cells in the samples. The absolute number of CD19<sup>+</sup> B cells actually underestimates the therapeutic efficacy of the treatment for elimination of CD20<sup>+</sup> B cells, however, since CD19 is expressed on B cell progenitor cells before expression of CD20.

#### **EXAMPLE 3**

Mice were injected five times over two months with one of four treatment protocols as follows:

human CD20 (44 aa fragment)-KLH plus QS1 human CD20 (44 aa fragment)-KLH human CD20 (44 aa fragment) plus QS21

KLH plus QS21

20

25

30

Blood was collected on week 9 for analysis by ELISA. Sera from the vaccinated mice were diluted 1:200 and incubated on BSA blocked plates coated with msCD20, huCD20, P190 or KLH. Secondary goat anti-mouse antibody conjugated to alkaline phosphatase was added, and the color change of p-nitrophenyl phosphate substrate was measured at 405 nm. The results are summarized in Figs. 7A-D. In most instances the peptide, carrier protein and adjuvant are all needed for optimal response, although some responses were detected using less than all of the components.

EXAMPLE 4

Mice were vaccinated according to the schedule of Example 3 using one of four treatment protocols: human CD20 (44 aa fragment)-KLH plus QS21 adjuvant, mouse

5

CD20 (44 amino acid fragment)-KLH plus QS21, P190 (irrelevant protein)-KLH +QS21 and KLH and QS21 alone. Mouse serum samples were evaluated by ELISA for the presence of antibodies reactive with msCD20, huCD20, P190 and KLH. The results are shown in Figs. 6A-D. Antibodies generated in mice after vaccination with human or mouse-derived CD20 fragments are specific for the peptides used, yet are capable of inducing immunity to the corresponding peptide from other species.

#### **EXAMPLE 5**

Mice were vaccinated five times over two months with huCD20 fragment-KLH conjugate with no adjuvant or in combination with one of three adjuvants: QS21, BCG or Alum. Serum samples from the vaccinated mice were tested by ELISA. The results are summarized in Figs. 8A-D. QS21 was found to be the most effective of those tested.

#### EXAMPLE<sub>6</sub>

To confirm the observations of Example 2, nucleated spleen cells were recovered by centrifugation in a density gradient from mice vaccinated with a CD20-KLH conjugate (human or mouse) in the presence of QS21 adjuvant. 1 X 106 cells from each mouse were incubated with 2  $\mu g$  of rat anti-mouse CD19 FITC-labeled antibody or with isotope-matched FITC labeled rat antibody. Cells were washed, fixed and analyzed with a Becton Dickinson FACScaliber cytometer. Figs 9A-C, D-F and G-I show the results for three exemplary mice of each vaccination group. The decrease in the peak reflecting levels of CD19 positive spleen cells in each of the mice is apparent.

*	<u></u>
_	9
٠,	0
F	a

Score (T <sub>1/2</sub> of Dissociation of Molecule Containing this Subsequence)	
Score ( Molect	1600 48 21.2 1600 60 89.4 28.7
HLA Molecule	Kd Kd A_0201 Kd Kd A_0201 A_0201
Peptide Sequence	NFIRAHTPYI FIRAHTPYI FLKMCSLNFI HFLKMRRLEL IYDCePSNSS LIQTSKPYV ELIQtSKPYV
Species	human human mouse mouse mouse

1 2

1

2

3

4

1

1


#### **CLAIMS**

1. A method for active vaccination against autologous cells expressing
transmembrane proteins comprising administering to a patient a vaccine composition
comprising at least an immunogenic portion of the extracellular domain of the
transmembrane protein, or a xenogeneic homolog thereof, coupled to or administered with an
carrier protein effective to break tolerance to the transmembrane protein and a
pharmaceutically acceptable adjuvant.
2. The method of claim 1, wherein the transmembrane protein is selected
from the group consisting of CD20. Her2-neu, VEGF receptor, epidermal growth factor

- 2. The method of claim 1, wherein the transmembrane protein is selected from the group consisting of CD20, Her2-neu, VEGF receptor, epidermal growth factor receptor, the CD19 molecule, interleukin-2-receptor, interleukin-4-receptor, and the P-glycoprotein.
  - 3. The method of claim 1, wherein the transmembrane protein is CD20.
- 4. The method of claim 1, wherein the vaccine composition comprises a peptide having the sequence given by Seq. ID No 1 or 2.
- 5. The method claim 1, wherein the carrier protein is keyhole limpet hemocyanin.
- 6. The method of claim 5, wherein the transmembrane protein is selected from the group consisting of CD20, Her2-neu, VEGF receptor, epidermal growth factor receptor, the CD19 molecule, interleukin-2-receptor, interleukin-4-receptor, and the P-glycoprotein.
  - 7. The method of claim 5, wherein the transmembrane protein is CD20.

- 8. The method of claim 7, wherein the vaccine composition comprises a peptide having the sequence given by Seq. ID No 1 or 2.
- 9. A method for active vaccination against B cells expressing CD20 comprising administering to a patient a vaccine composition comprising at least an immunogenic portion of the extracellular domain of CD20, or a xenogeneic homolog thereof, coupled to or administered with an carrier protein effective to break tolerance to the transmembrane protein and a pharmaceutically acceptable adjuvant.
- 10. The method claim 9, wherein the carrier protein is keyhole limpet hemocyanin.
- 11. The method of claim 9, wherein the vaccine composition comprises a peptide having the sequence given by Seq. ID No 1 or 2.
- 12. A method for treatment of B cell non-Hodgkin's lymphoma, comprising administering to a patient suffering from B cell non-Hodgkin's lymphoma a vaccine composition comprising at least an immunogenic portion of the extracellular domain of CD20, or a xenogeneic homolog thereof, coupled to or administered with an carrier protein effective to break tolerance to the transmembrane protein and a pharmaceutically acceptable adjuvant.
- 13. A vaccine composition comprising at least an immunogenic portion of the extracellular domain of the transmembrane protein, or a xenogeneic homolog thereof, coupled to or administered with an carrier protein effective to break tolerance to the transmembrane protein and a pharmaceutically acceptable adjuvant.
- 14. The composition of claim 13, wherein the transmembrane protein is selected from the group consisting of CD20, Her2-neu, VEGF receptor, epidermal growth

3	factor receptor, the CD19 molecule, interleukin-2-receptor, interleukin-4-receptor, and the
4	P-glycoprotein.
1	15. The composition of claim 13, wherein the transmembrane protein is
2	CD20.
1	16. The composition of claim 15, wherein the vaccine composition
2	comprises a peptide having the sequence given by Seq. ID No 1 or 2.
400 km km 2 2 fm and 1 1	17. The composition of claim 13, wherein the carrier protein is keyhole
2	limpet hemocyanin.
1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1	18. The composition of claim 17, wherein the transmembrane protein is
<b>£</b> 2	selected from the group consisting of CD20, Her2-neu, VEGF receptor, epidermal growth
3	factor receptor, the CD19 molecule, interleukin-2-receptor, interleukin-4-receptor, and the P-
3 4	glycoprotein.
1	19. The composition of claim 17, wherein the transmembrane protein is
2	CD20.
1	20. The composition of claim 19, wherein the vaccine composition
2	comprises a peptide having the sequence given by Seq. ID No 1 or 2.








FIG. 1B



TOTAL EVENTS: 10000

			~
MARKER	LEFT, RIG	HT EVENTS	% TOTAL
ALL	1, 99		100.00
M1	3,	7525	75.25
M2	1	3 1201	12.01
IVI ←	' 7	<del>-</del>	

FIG. 2A




TOTAL EVENTS: 10000

MARKER	LEFT, RIGHT	<b>EVENTS</b>	% TOTAL
ALL M1	1, 9910 3, 12	10000 7327	100.00 73.27 1 4.54
M2	1, 4	1454	14.54


FIG. 2B

09/87/970




3/9

FIG. 3



**SUBSTITUTE SHEET (RULE 26)** 







FIG. 6A

FIG. 6B






FIG. 6C

FIG. 6D




FIG. 7A




FIG. 7B




FIG. 7C




FIG. 7D




FIG. 8A




FIG. 8B




FIG. 8C






FIG. 8D







#### SEQUENCE LISTING

```
<110> Agus, David B.
        Scheinberg, David
        Zelenetz, Andrew D.
        Roberts, Wenay
  <120> Compositions and Methods for Active Vaccination
  <130> MSKP039US
  <140>
 <141>
 <150> 60/084,370
 <151> 1998-05-08
 <160> 6
 <170> PatentIn Ver. 2.0
<210> 1
 <211> 44
 <212> PRT
 <213> HUMAN
 <220>
 <223> human CD20 fragment
 <400> 1
 Lys Ile Sor His Phe Lou Lys Met Glu Ser Lou Asn Phe Ile Arg Ala
                      10
        5
His Thr Pro Tyr Ile Asn Ile Tyr Asn Cys Glu Pro Ala Asn Pro Ser
                                     30
           20 25
Glu Lys Asn Ser Pro Ser Thr Gln Tyr Cys Tyr Ser
        35
<210> 2
<211> 44
<212> PRT
<213> Murine
<220>
<223> murine CD20 fragment
```

```
<400> 2
   Thr Leu Ser His Phe Lou Lys Met Arg Arg Leu Glu Leu Ile Gln Thr
                                       10
   Ser Lys Pro Ty: vel Asp Ile Ty: Asp Cys Glu Pro Ser Ash Ser Ser
                                    25
  Glu Lys Asn Ser Pro Ser Thr Gln Tyr Cys Asn Ser
           35
                              40
  <210> 3
  <211> 79
  <212> PRT
  <213> HUMAN
 <220>
 <223> exon VI of numan CD20 gene
 <400> 3
 Val Lys Gly Lys Met Ile Met Asn Ser Lou Ser Leu Phe Ala Ala Ile
                   5
                                      10
 Ser Gly Met Ile Leu Ser Ile Met Asp Ile Leu Asn Ile Lys Ile Ser
             20
                                  25
                                                       30
 His Phc Lou Lys His Glu Ser Leu Asn Phe Ile Arg Ala His Thr Pro
                              40
 Tyr Ile Asn Ile Tyr Asn Cys Glu Pro Ala Asn Pro Ser Glu Lys Asn
      50
                         55
 Ser Pro Ser Thr Glm Tyr Cys Tyr Ser Ile Gln Ser Leu Phe Leu
 65
                     70
<210> 4
<211> 237
<212> DNA
<213> HUMAN
<220>
<223> exon VI of human CD20 gene
<400> 4
gtcaaaggaa aaatgataat gaattcattg agoototttg otgocattto tggaatgatt 60
ctttcaatca tggacttoot taatattaaa atttcccatt ttttaaaaat ggagagtctg 120
aattttatta gageteasac accatatatt aacatataca actgtgaacc agetaateec 180
```

tetgagaaca actecceate tacceaatac tgttacagea tacaatetet gttettg	237
<210> 5 <211> 1255 <212> PRT <213> HUMAN	
(213) II3.04(	
<220> <223> numan Her2	
<400> 5	
Met Glu Leu Ala Ala Leu Cys Arg Trp Gly Leu Leu Leu Leu 1 5 16 15	
Pro Pro Cly Ala Ala Ser Thr Gln Val Cys Thr Gly Thr Asp Met Lys 25 30	
Lou Arg Leu Pro Ala Ser Pro Glu Thr His Leu Asp Met Leu Arg His 35 40 45	
Leu Tyr Gln Gly Cys Gln Val Val Gln Gly Asn Leu Glu Leu Thr Tyr 50 55 60	
Leu Pro Thr Asn Ala Ser Leu Ser Phe Leu Gln Asp Ile Gln Glu Val 65 70 75 80	
Gln Gly Tyr Val Leu Ile Ala His Asn Gln Val Arg Gln Val Pro Leu 85 90 95	
Gln Arg Lou Arg Ile Val Arg Gly Thr Gln Leu Phe Glu Asp Asn Tyr 100 105 110	
Ala Leu Ala Val Leu Asp Asn Gly Asp Pro Leu Asn Asn Thr Thr Pro 115 120 125	
Val Thr Gly Ala Ser Pro Gly Gly Leu Arg Glu Leu Gln Leu Arg Ser 130 135 140	
Leu Thr Glu Ile Leu Lys Gly Gly Val Leu Ile Gln Arg Asn Pro Gln 145 150 155 160	
Leu Cys Tyr Gln Asp Thr Ile Leu Trp Lys Asp Ile Phe His Lys Asn 135 170 175	
Asn Glr Leu Ala Leu Thr Leu Ile Asp Thr Asn Arg Ser Arg Ala Cys 180 185 190	

His Pro Cys Ser Pro Met Cys Lys Gly Ser Arg Cys Trp Gly Glu Ser

195	200	205
Ser Glu Asp Cys Gln Ser	Leu Thr Arg Thr	Val Cys Ala Gly Gly Cys
210	215	220
Ala Arg Cys Lys Gly Pro 225 230		Cys Cys His Glu Gln Cys 235 240
Ala Ala Gly Cys Thr Gly	Pro Lys His Sor .	Asp Cys Leu Ala Cys Leu
245	250	255
His Phe Asn His Ser Gly	Ile Cys Glu Leu F	His Cys Pro Ala Leu Val
260	265	270
Thr Tyr Ash Thr Asp Thr : 275	Phe Glu Sor Met F 280	Pro Asn Pro Glu Gly Arg 285
Tyr Thr Pne Gly Ala Scr (	Cys Val inm Ala C	ys Pro Tyr Asn Tyr Leu
290	295	300
Ser Thr Asp Val Gly Ser 0 305 310	-	ys Pro Leu His Asn Gln 15 320
Glu Val Thr Ala Glu Asp G	Gly Thr Gln Arg Cy	ys Glu Lys Cys Ser Lys
325	330	335
Pro Cys Ala Arg Val Cys T	yr Gly Lea Gly Ma	et Glu His Leu Arg Glu
340	345	350
Val Arg Ala Val Thr Ser A	la Asn Ile Gln Gl	u Pnc Ala Gly Cys Lys
355	360	365
Lys Ile Phe Gly Ser Leu Al	la Phe Leu Pro Gl 75	u Ser Pae Asp Gly Asp 380
Pro Ala Ser Asn Thr Ala Pr 385 390	co Leu Gln Pro Gl 39	
Glu Thr Leu Glu Glu Ile Th	nr Gly Tyr Leu Ty:	r Ile Ser Ala Trp Pro
405	410	415
Asp Ser Leu Pro Asp Leu Se	r Val Phe Gln Asr	n Leu Glm Val Ile Arg
420	425	430
Gly Arg Ile Leu His Asn Gly	y Ala Tyr Ser Leu	Thr Leu Gln Gly Leu
435	440	445
Gly lie Ser Trp Leu Gly Leu	u Arg Ser Leu Arg	Glu Leu Gly Ser Gly

450	455	460	
Leu Ala Leu Ile H 465	is His Ash Thr His	Leu Cys Phe Val :	His Thr Val
	eu Pne Arg Asn Pro 3	His Gln Ala Leu I 490	Lou His Thr 495
Ala Asn Arg Pro Gl	iu Asp Glu Cys Val (		Ala Cys His
500	505		510
Gln Leu Cys Ala Ar	rg <b>C</b> ly His Cys Trp G	Gly Pro Gly Pro T	hr Gln Cys
515	520	525	
Val Asn Cys Ser Gl	n Phe Let Arg Gly G	iln Glu Cys Val G	lu Glu Cys
530	535	540	
Arg Val Leu Glm Gl;	y Leu Pro Arg Glu T	yr Val Asn Ala Ar	rg His Cys
545	550	555	560
Lou Pro Cys His Pro	o Glu Cys Gln Pro G	ln Asn Gly Ser Va	al Thr Cys
565	5	70	575
Phe Gly Pro Glu Ala	a Asp Gln Cys Val Al	la Cys Ala His Ty	
580	585	59	
Pro Pro Phe Cys Val	Ala Arg Cys Pro Se	er Gly Val Lys Pro	o Asp Leu
595	600	605	
Ser Tyr Met Pro Ile 610	Trp Lys Phe Pro As	p Glu Glu Gly Ala 620	a Cys Gln
Pro Cys Pro Ile Asn	Cys Thr His Ser Cys	s Val Asp Leu Asp	Asp Lys
625		635	640
Gly Cys Pro Ala Glu	Gln Arg Ala Ser Pro		e Ile Ser
645	650		655
Ala Val Val Gly Ile	Leu Leu Val Val Val	l Leu Gly Val Val	
660	665	670	
Ile Leu Ile Lys Arg	Arg Gln Gln Lys Ile	Arg Lys Tyr Thr	Met Arg
675	680	685	
Arg Leu Leu Gln Jlu 690	Thr Clu Leu Val Glu 695	Pro Lou Thr Pro	Ser Gly
Ala Met Pro Asn Gln A	Ala Gln Met Arg Ile	Leu Lys Glu Thr	Glu Leu

WO 99/57981 6 PCT/US99/10065

705	710	715 720
Arg Lys Val Lys Val L 725	cu Gly Ser Gly Ala 730	Phe Gly Thr Val Tyr Lys 735
Gly lle Trp lle Pro A.	sp Gly Glu Asn Val 1 745	Lys Ile Pro Val Ala Ile 750
Lys Val Leu Arg Glu As 755	sn Thr Ser Pro Lys ? 760	Ala Asn Lys Glu Ile Leu 765
Asp Glu Ala Tyr Val Mc 770	et Ala Gly Val Gly S 775	Ser Pro Tyr Val Ser Arg 780
Lou Leu Gly Ile Cys Lo 785 79		in Leu Val Thr Gln Leu 95 800
Met Pro Tyr Gly Cys Le 805	eu Leu Asp His Val A 810	rg Clu Asn Arg Gly Arg 815
Leu Gly Ser Glm Asp Let 820	u Leu Asn Trp Cys Mo 825	et Gln Ile Ala Lys Gly 830
Met Ser Tyr Leu Glu Asg 835	p Val Arg Leu Val Hi 840	is Arg Asp Leu Ala Ala 845
Arg Asn Val Leu Val Lys 850	s Ser Pro Asn His Va 855	l Lys Ile Thr Asp Phe 860
Gly Leu Ala Arg Leu Leu 965 870		
Gly Gly Lys Val Pro Ile 885	Lys Trp Met Ala Le 890	u Glu Ser Ile Leu Arg 895
Arg Arg Phe Thr His Gln 900	Ser Asp Val Trp Ser 905	r Tyr Gly Val Thr Val 910
Trp Glu Lou Met Thr Phe 915	Gly Ala Lys Pro Tyr 920	Asp Gly Ile Pro Ala
Arg Glu Ile Pro Asp Leu 930	Leu Glu Lys Gly Glu 935	Arg Leu Pro Gln Pro
Pro Ile Cys Thr Ile Asp 945 950	Val Tyr Met Ile Met 955	
Ile Asp Ser Glu Cys Arg	Pro Arg Phe Arg Glu	Leu Val Ser Glu Phe

965	970	975
Ser Arg Met Ala Arg Asp Pro Gln Arg 980 985	Phe Val Val Ile Gln	
Asp Leu Gly Pro Ala Ser Pro Leu Asp 995 1000	Ser Thr Phe Tyr Arg 1005	Ser Leu
Leu Glu Asp Asp Asp Met Gly Asp Leu 1 1010 1015	Val Asp Ala Glu Glu 1020	Tyr Leu
Val Pro Gln Gln Gly Phe Phe Cys Pro A 1025 1030	Asp Pro Ala Pro Gly 1035	Ala Gly 1040
Gly Not Val His His Arg His Arg Ser S 1045		Gly Gly .055
Gly Asp Leu Thr Leu Gly Leu Glu Pro S 1060 1065	er Glu Glu Glu Ala 1070	Pro Arg
Ser Pro Leu Ala Pro Ser Glu Cly Ala G 1075 1080	ly Ser Asp Val Phe 1085	Asp Gly
Asp Leu Gly Met Gly Ala Ala Lys Gly Lo	eu Gln Ser Leu Pro ' 1100	Thr His
Asp Pro Ser Pro Leu Gln Arg Tyr Ser Gl 1105 1110	Hu Asp Pro Thr Val 1 1115	Pro Leu 1120
Pro Ser Glu Thr Asp Gly Tyr Val Aia Pr 1125 113		Pro Gln .35
Pro Glu Tyr Val Asn Gln Pro Asp Val Ar 1140 1145	g Pro Gln Pro Pro S 1150	er Pro
Arg Glu Gly Pro Leu Pro Ala Ala Arg Pr 1155 1160	o Ala Gly Ala Thr L 1165	eu Glu
Arg Pro Lys Thr Leu Ser Pro Gly Lys As:	n Gly Val Val Lys A: 1180	sp Val
Phe Ala Phe Gly Gly Ala Val Glu Asn Pro	o Glu Tyr Lou Thr Pi 1195	co Gln 1200
Gly Gly Ala Ala Pro Gln Pro His Pro Pro 1205 1210		
Phe Asp Asn Leu Tyr Tyr Trp Asp Gln Asp	Pro Pro Glu Arg Gl	y Ala

WO 99/57981 PCT/US99/10065

8

1220 1225 1230

Pro Pro Ser Thr Phe Lys Gly Thr Pro Thr Ala Glu Asn Pro Glu Tyr 1235 1240 1245

Leu Gly Leu Asp Val Pro Val 1250 1255

<210> 6

<211> 1210

<212> PRT

<213> HUMAN

<220>

<223> numan EGFR

<400> 6

Met Arg Pro Ser Gly Thr Ala Gly Ala Ala Leu Leu Ala Leu Leu Ala 1 5 10 15

Ala Leu Cys Pro Ala Ser Arg Ala Leu Glu Glu Lys Lys Val Cys Gln \$20\$  $$25\,_{-}$$  30

Gly Thr Ser Asn Lys Leu Thr Gln Leu Gly Thr Phe Glu Asp His Phe 35 40 45

Leu Ser Leu Gln Arg Met Phe Asn Asn Cys Glu Val Val Leu Gly Asn 50 55 60

Leu Glu Ile Thr Tyr Val Gln Arg Asn Tyr Asp Leu Ser Phe Leu Lys
65 70 75 80

Thr IIc Gln Glu Val Ala Gly Tyr Val Leu IIc Ala Leu Asn Thr Val
85 90 95

Glu Arg Ile Pro Leu Glu Asn Leu Gln Ile Ile Arg Gly Asn Met Tyr 100 105 110

Tyr Glu Asn Ser Tyr Ala Leu Ala Val Leu Ser Aon Tyr Asp Ala Asn 115 \$120\$ 125

Lys Thr Gly Leu Lys Glu Leu Pro Met Arg Asn Leu Gln Glu Ile Leu 130 135 140

Ser Ile Gln Trp Arg Asp Ile Val Ser Scr Asp Phe Leu Se 165 170	r Asn Met 175
Ser Met Asp Phe Gln Asn His Leu Gly Ser Cys Gln Lys Cys 180 185 190	
Ser Cys Pro Asn Gly Ser Cys Trp Gly Ala Gly Glu Glu Asn 195 200 205	n Cys Gln
Lys Leu Thr Lys Ile Ile Cys Ala Gin Gln Cys Ser Gly Arg 210 215 220	Cys Arg
Gly Lys Ser Pro Ser Asp Cys Cys His Asn Gln Cys Ala Ala 225 230 235	Gly Cys 240
Thr Gly Pro Arg Glu Ser Asp Cys Leu Val Cys Arg Lys Phe 245 250	Arg Asp 255
Glu Ala Thr Cys Lys Asp Thr Cys Pro Pro Leu Met Leu Tyr 260 265 270	Asn Pro
Thr Tnr Tyr Gln Met Asp Val Asn Pro Glu Gly Lys Tyr Scr 275 280 285	Phe Gly
Ala Thr Cys Val Lys Lys Cys Pro Arg Asn Tyr Val Val Thr 290 295 300	Asp His
Gly Ser Cys Val Arg Ala Cys Gly Ala Asp Ser Tyr Glu Met 305 310 315	Glu Glu 320
Asp Gly Val Arg Lys Cys Lys Lys Cys Glu Gly Pro Cys Arg 325 330	Lys Val 335
Cys Asn Gly Ile Gly Ilo Gly Glu Phe Lys Asp Ser Lou Ser 340 345 350	Ile Asn
Ala Thr Asn Ile Lys His Phe Lys Asn Cys Thr Ser Ile Ser (	Gly Asp
Leu His Ile Leu Pro Val Ala Phe Arg Gly Asp Ser Phe Thr F 370 375 380	His Thr
Pro Pro Leu Asp Pro Gln Glu Leu Asp Ile Leu Lyc Thr Val L 385 390 395	ys Glu 400
Ile Thr Gly Phe Leu Leu Ile Gln Ala Trp Pro Glu Asn Arg T	hr Asp

415

Le	u H	15 A		Phe ( 120	Glu i	Asn	Leu	Glu	11c		e Arg	; Gly	· Arg	Tnr 430	Lys	Gln
Hit	s G		iln I	he S	Ser I	œu i		Val 440	Va!	Ser	t ned	ı Asn	Ile 445		Ser	Leu
Gly	y Lo 45		rg S	er I	cu I		Glu 155	Ile	Ser	Asp	Gly	Asp 460	Val	Ile	Ile	Ser
Gly 465		sn L	ys A	sn l		ys T 70	yr .	9la	Asn	Thr	Ile 475	Asn	Trp	Lys	Lys	Leu 480
Phe	: Gl	y Tì	nr Se		ly G 35	ln L	ys T	Chr	Lys	Ile 490	Ile	Ser	Asn	Arg	Gly 495	Glu
Asn	Se	r СУ	's Ly 50		a Ti	ır G	ly C		Val 505	Суз	His	Ala		Cys 510	Ser	Pro
Glu	Gl	у Су 51		p Gi	y Pr	o Gi		ro 2 20	Arg .	Asp	Суѕ	Val	Ser 525	Cys <i>I</i>	Arg	Asn
Val	Sei 530		g Gl	y Ar	g Gl	u Cy 53		al A	Asp 1	Lys ·		Lys I 540	Leu I	Leu (	Glu	Gly
Glu 545	Pro	Ar	g Gl	u Ph	e Va 55		u As	sn S	er (		Cys : 555	ĭle 0	Gln (	Cys H		Pro 660
Glu	Суз	Let	ı Pro	565		a Me	t As	in I		hr (	Cys 1	Thr G	ly A		ly F 75	ro'ro
Asp A	Asn	Суа	: Ilc 580		суз	al.	a Hı		yr I 85	le A	sp C	Sly P		15 C	ys V	al
Lys 1	Thr	Cys 595	Pro	Ala	Gly	· Val	l Me 60		ly G	lu A	an A		hr L	eu Va	al T	rp
Lys T	Tyr 510	Ala	Asp	Ala	Gly	His 615		l Cy	/s H:	ıs L		ys H: 20	ıs Pı	ro As	sn C	ys
Thr T 625	,Àr	Gly	Cys	Tar	Gly 630	Pro	Gly	/ Le	u Gl		ly Cy 35	ys Pr	to Th	nr As	n Gl 64	
Pro L	ys	Ile	Pro	Ser 545	Ile	Ala	Thr	Gl	у Ма 65		al GJ	ly Al	a Le	u Le 65		ıu
Leu Lo	eu '	Val	Val 660	Ala	Leu	Gly	Ile	G1 <sub>3</sub>		u Ph	ne Mo	et Ar	g Ar	_	g Hı	ន

Ile		Arg : 675	Lys A	Arg T	Thr I		Arg 680	Arg	Leu	Lou	Gln	Glu 685	Arg	Glu	Leu
Val (	Glu I 690	Pro I	icu I	hr P		Ser (	Cly	Glu	Ala	₽ro	Asn 700	Gln	Ala	Leu	Leu
Arg 3	Ile I	.eu L	ys G		hr G 10	lu I	?he	Lys	Lys	Ile 715	Lys i	Val	Leu	Gly	Ser 720
Gly A	ala P	he G		nr V. 25	al T	yr L	ys		Leu 730	Trp :	ile E	Pro	Glu	Gly 735	Glu
Lys V	al L		le P: 40	ro Va	-1 A	la I		Lys 745	Glu	Leu i	\rg C		Ala 750	Thr	Ser
Pro L		la A: 55	sn Ly	rs Gl	u II		eu <i>I</i> 60	lsp (	Glu <i>i</i>	Ala T		al 1 65	Met	Ala	Ser
Val As	sp As 70	sn Pr	co Hi	s Va	1 Cy 77		rg I	eu I	Leu (	_	le C; 80	ys I	Jeu '	Thr	Ser
Thr Va	al Gl	n Le	u Il	e Th 79		n Le	eu M	ot F		he G 95	ly C	/s L	.cu I		Asp 300
Tyr Va	ıl Ar	g Gl	u Ale 805		s As	p As	n I		ly S	er Gl	ln Ty	r L		eu <i>P</i>	sn
Trp Cy	s Va	1 Gl:		e Alá	ı Ly:	s Gl		et A: 25	sn T	yr Le	eu Gl		sp A 30	rg A	rg
Leu Va	835		) Asp	Leu	Ala	840		rg As	sn Va	al Le	u Va 84		ys T	nr P	ro
Gln His		Lys	: Ile	Thr	Asp 855		e Gl	y Le	eu Al	a Ly 86		ı Le	eu G	ly A	la
Glu Glu 865	ı Lys	Glu	Tyr	His 870	Ala	Glu	ı Gl	y Gl	y Ly 87		l Pro	o Il	e Ly	/s T1 88	
Met Ala	Leu	Glu	Ser 385	Ilc	Leu	His	Ar	g Il 89		r Thi	: His	Gl	n Se 89		īĐ
Val Trp	Ser	Tyr 900	Gly	Val	Thr	Val	Tr:		u Le	ı Met	Thr	Ph:		y Se	r
Lys Pro	Tyr 915	Asp	Gly	Ile	Pro	Ala 920	Ser	Glu	ı Ile	e Ser	Ser 925	Ile	e Le	u Gl	ս

- Lys Gly Glu Arg Leu Pro Gln Pro Pro Ile Cys Thr Ile Asp Val Tyr 920 935 940
- Met Ile Met Val Lys Cys Trp Met Ile Asp Ala Asp Sar Arg Pro Lys 945 950 955 960
- Phe Arg Glu Leu Ile Ile Glu Phe Ser Lys Met Ala Arg Asp Pro Gln 965 970 975
- Arg Tyr Lou Val Ile Gln Gly Asp Glu Arg Met His Leu Pro Ser Pro 980 985 990
- Thr Asp Ser Ash Phe Tyr Arg Ala Leu Met Asp Glu Glu Asp Met Asp 995 1000 1005
- Asp Val Val Asp Ala Asp Glu Tyr Lou Ilo Pro Gln Gln Gly Phe Phe 1010 1015 1020
- Ser Ser Pro Ser Thr Ser Arg Thr Pro Leu Leu Ser Ser Leu Ser Ala 1025 1030 1035 1040
- Thr Ser Asn Asn Ser Thr Val Ala Cys Ile Asp Arg Asn Gly Leu Gln 1045 1050 1055
- Ser Cys Pro Ile Lys Glu Asp Ser Phe Leu Gln Arg Tyr Ser Ser Asp 1060 1065 1070
- Pro Thr Gly Ala Leu Thr Glu Asp Ser Ile Asp Asp Thr Phe Leu Pro 1075 1080 1085
- Val Pro Glu Tyr Ile Asn Gln Ser Val Pro Lys Arg Pro Ala Gly Ser 1090 1095 1100
- Val Gln Asn Pro Val Tyr His Asn Gln Pro Leu Asn Pro Ala Pro Ser 1105 1110 1115 1120
- Arg Asp Pro His Tyr Gln Asp Pro His Ser Thr Ala Val Gly Asn Pro 1125 1130 1135
- Glu Tyr Leu Asn Thr Val Gln Pro Thr Cys Val Asn Ser Thr Phe Asp 1140 1145 1150
- Ser Pro Ala His Trp Ala Gln Lys Gly Scr His Gln Ile Ser Leu Asp 1155 1160 1165
- Ash Pro Asp Tyr Gln Gln Asp Phe Phe Pro Lys Glu Ala Lys Pro Ash 1170 1175 1180

Gly Ile Phe Lys Gly Ser Thr Ala Glu Asn Ala Glu Tyr Leu Arg Val 1185 1190 1195 1200

Ala Pro Gln Ser Ser Glu Phe Ile Gly Ala 1205 1210

OPPEDAHL & LARSON

FILE NO. MSKP039 INVENTOR . Agus, et al

Claim for Priority

I hereby claim foreign priority benefits under 35 U.S.C. § 119 (a)-(d) or 365(b) of any foreign application(s) for patent or inventor's certificate, or 365(a) of any PCT international application which designated at least one country other than the United States of America, listed below and have also identified below any foreign applications for patent or inventor's certificate, or of any PCT international application having a filing date before that of the application on which priority is claimed.

		(day/month/year)	CLAIMED	COPY ATTACHED
	· <u>-</u>		YES[]NO[]	YES[]NO[]
ON(S), IF ANY, FILED MO	ORE THAN 12 MONTH	IS (6 MONTHS FOR D	DESIGN) PRIOR T	O SAID
APPLICATION NO.	DATE OF FILING (day/month/year)	DATE OF ISSUE (day/month/year)		
		APPLICATION NO. DATE OF FILING	APPLICATION NO. DATE OF FILING DATE OF ISSUE	ON(S), IF ANY, FILED MORE THAN 12 MONTHS (6 MONTHS FOR DESIGN) PRIOR TO APPLICATION NO. DATE OF FILING DATE OF ISSUE

l nereby claim the benefit under 35 U.S.C § 119(e) of any United States provisional application(s) listed below.

60/084,870	08 May 1998	
(application number)	(filing date)	·.
		•
(application number)	(filing date)	

I hereby declare that all statements made herein of my own knowledge are true and that all statements made on information and belief are believed to be true; and further that these statements are made with the knowledge that willful false statements and the like so made are punishable by fine or imprisonment or both, under Section 1001 of Title 18 of the United States Code and that such willful false statements may jeopardize the validity of the application or any patent issued thereon.

NAME OF SOLE OR FIRST INVENTOR	LAST NAME AGUS	FIRST NAME	MIDDLE NAME B.
RESIDENCE & CITIZENSHIP	CITY OF RESIDENCE  Brocklyn Beverly Hills PM	STATE OR COUNTRY OF RESIDENCE OF CA	COUNTRY OF CITIZENSHIP US
POST OFFICE ADDRESS Pierrepont Street 521 North (		CITY Brooklyn PA Bevery Hills	STATE/COUNTRY ZIP CODE NY 10021 PA CA 9 02 1 0
DATE () o	0	SIGNATURE	

[X] Signature for additional joint inventor attached. Number of Pages 1. [] Signature by Administrator(trix) or legal representative for deceased or

incapacitated inventor. Number of Pages \_\_\_.

[] Signature for inventor who refuses to sign or cannot be reached by person authorized under 37 CFR § 1.47. Number of Pages \_\_\_.

PAGE 03/04

FILE NO. MSKP039 INVENTOR . Agus, et al

NAME OF SECOND INVENTOR	LAST NAME SCHEINBERG	FIRST NAME DAVID	MIDDLE NAME
RESIDENCE & CITY OF RESIDENCE NEW YORK		STATE OR COUNTRY OF RESIDENCE NY	COUNTRY OF CITIZENSHIP US
POST OFFICE ADDRE 325 Central Park V		CITY New York	STATE/COUNTRY ZIP CODE NY 10025
DATE \	Co	SIGNATURE	<del>)</del>
NAME OF THIRD INVENTOR	LAST NAME ROBERTS	FIRST NAME WENDY	MIDDLE NAME
RESIDENCE & CITIZENSHIP	CITY OF RESIDENCE NEW YORK	STATE OR COUNTRY OF RESIDENCE NY	COUNTRY OF CITIZENSHIP US
POST OFFICE ADDRE 1233 York Avenue	iss , 303 East 715+	CITY New York	STATE/COUNTRY ZIP CODE NY 10021
DATE II/I/DU	)	SIGNATURE WHUME PHU	
NAME OF FOURTH INVENTOR	LAST NAME ZELENETZ	FIRST NAME ANDREW	MIDDLE NAME D.
RESIDENCE & CITIZENSHIP	CITY OF RESIDENCE LARCHMONT	STATE OR COUNTRY OF RESIDENCE NY	COUNTRY OF CITIZENSHIP US
POST OFFICE ADDRE		CITY Larchmont	STATE/COUNTRY ZIP CODE NY 10538
DATE 11/6/2003	` >	SIGNATURE Ann telem	

FILE NO. MSKP039 INVENTOR . Agus, et al

# COMBINED DECLARATION AND POWER OF ATTORNEY

As a below named inventor, I hereby declare that:

My citizenship, residence and post office address are as listed below next to my name.

-					
l believ which a	e I am the original a patent is sought o	, first and [ ] sole/[x on the invention er	<ul> <li>i) joint inventor of the sultitled: Compositions and</li> </ul>	bject matter whi Methods for Ac	ch is claimed and for tive Vaccination
the spe	cification of which				
(a)[]	is attached hereto	<b>D</b> .			
(b)[]	was filed on	as	Application Serial No.	,	and was amended
(c) [X]	was described ar and amended on	nd claimed in Inter	national Application No	PCT/US99/1000	65 filed on <u>May 7, 1999</u>
includi	ng the claims, as a ation which is mate	reviewed and und imended by any ar erial to the patenta	edgment of Duty of Disc erstood the content of the mendment referred to ab bility of the subject matte Regulations § 1.56(a).	ne above identifi nove. I acknowle	BUMB file anth to algologe
365(c) insofar States	of any PCT internation or PCT internation whedge the duty to en the filing date or	ational application tter of each of the nal application in the disclose material i	on and the national or P	n is not disclose the first paragrap 37 CFR § 1.56 CT internationa	d in the prior United on of 35 U.S.C. § 112, I which became available I filing date of this
(Applica	tion Serial No.)	(Filing Date)	(Status)(patented,pending	g,abandoned)	(Patent No. if applicable)
(Applica	tion Serial No.)	(Filing Date)	(Status)(patented,pending	g,abandoned)	(Patent No. if applicable)
			Power of Attorney		
Nancy	J. Parsons, PTO	Reg. No. 40,364 o Center 2 <sup>nd</sup> Floor 2	NO. 32,746, Marina T. I f the firm of OPPEDAHL 56 Dillon Ridge Rd., Dillo in the Patent and Trade	on. CO 80435 a	is attorneys to prosecute
l.	CORRESPONDENCE  O21121	40	DIRECT TELEP OPPEDAHL & L (970) 468-6600	HONE CALLS TO: ARSON LLP	·

PATENT TRADEHARK OFFICE

FILE NO. MSKP039 INVENTOR . Agus, et al

**Claim for Priority** 

I hereby claim foreign priority benefits under 35 U.S.C. § 119 (a)-(d) or 365(b) of any foreign application(s) for patent or inventor's certificate, or 365(a) of any PCT international application which designated at least one country other than the United States of America, listed below and have also identified below any foreign applications for patent or inventor's certificate, or of any PCT international application having a filing date before that of the application on which priority is claimed.

EARLIEST FOREIGN APPLICATION	N APPLICATION(S), FILED	WITHIN TWELVE MO	NTHS (6 MONTHS F	OR DESIGN) PRIC	OR TO SAID
COUNTRY	APPLICATION NO.	DATE OF FILING (day/month/year)	DATE OF ISSUE (day/month/year)	PRIORITY CLAIMED	CERTIFIED COPY ATTACHED
				YES[] NO[]	YES[]NO[]
FOREIGN APPLICAT	TION(S), IF ANY, FILED MO	ORE THAN 12 MONTH	HS (6 MONTHS FOR I	DESIGN) PRIOR T	O SAID
COUNTRY	APPLICATION NO.	DATE OF FILING (day/month/year)	DATE OF ISSUE (day/month/year)		
		Provisional App	olication		
•	benefit under 35 U.S.	C § 119(e) of any	United States pro	visional applica	ation(s) listed
below.		00 M-	1000		
60/084,870			y 1998		
(application number	er)	(filing d	ate)		
(application number	er)	(filing d	ate)		

I hereby declare that all statements made herein of my own knowledge are true and that all statements made on information and belief are believed to be true; and further that these statements are made with the knowledge that willful false statements and the like so made are punishable by fine or imprisonment or both, under Section 1001 of Title 18 of the United States Code and that such willful false statements may jeopardize the validity of the application or any patent issued thereon.

NAME OF SOLE OR FIRST INVENTOR	LAST NAME AGUS	FIRST NAME David	MIDDLE NAME B.
RESIDENCE & CITIZENSHIP	CITY OF RESIDENCE Brooklyn	STATE OR COUNTRY OF RESIDENCE NY	COUNTRY OF CITIZENSHIP US
POST OFFICE ADDRESS 9 Pierrepont Street		CITY Brooklyn	STATE/COUNTRY ZIP CODE NY 10021
DATE		SIGNATURE	

[X] Signature for additional joint inventor attached. Number of Pages 1.

[] Signature by Administrator(trix) or legal representative for deceased or incapacitated inventor. Number of Pages \_\_\_.

[] Signature for inventor who refuses to sign or cannot be reached by person authorized under 37 CFR § 1.47. Number of Pages \_\_\_.

OPPEDAHL & LARSON

FILE NO. <u>MSKP039</u> INVENTOR . Agus, et al

NAME OF SECOND INVENTOR	LAST NAME SCHEINBERG	FIRST NAME DAVID	MIDDLE NAME
RESIDENCE & CITIZENSHIP	CITY OF RESIDENCE NEW YORK	STATE OR COUNTRY OF RESIDENCE NY	COUNTRY OF CITIZENSHIP US
POST OFFICE ADDRESS 325 Central Park West		CITY New York	STATE/COUNTRY ZIP CODE NY 10025
DATE		SIGNATURE	
NAME OF THIRD	LAST NAME ROBERTS	FIRST NAME WENDY	MIDDLE NAME
RESIDENCE & CITIZENSHIP	CITY OF RESIDENCE NEW YORK	STATE OR COUNTRY OF RESIDENCE NY	COUNTRY OF CITIZENSHIP US
POST OFFICE ADDRESS 1233 York Avenue		CITY New York	STATE/COUNTRY ZIP CODE NY 10021
DATE		SIGNATURE	
NAME OF FOURTH INVENTOR	LAST.NAME ZELENETZ	FIRST NAME ANDREW	MIDDLE NAME D.
RESIDENCE & CITIZENSHIP	CITY OF RESIDENCE LARCHMONT	STATE OR COUNTRY OF RESIDENCE NY	COUNTRY OF CITIZENSHIP US
POST OFFICE ADDR 31 Mohegan Road		CITY Larchmont	STATE/COUNTRY ZIP CODE NY 10538
DATE		SIGNATURE	·

**OPPEDAHL & LARSON** 

FILE NO. MSKP039 INVENTOR . Agus, et al

## COMBINED DECLARATION AND POWER OF ATTORNEY

As a below named inventor, I hereby declare that:

My citizenship, residence and post office address are as listed below next to my name.

I believe I am the original, first and [] sole/[x] joint inventor of the subject matter which is claimed and for which a patent is sought on the invention entitled: Compositions and Methods for Active Vaccination

	patern to cought		THE COMPOSITOR OF THE WILLIAM OF THE	THOUTO TOOMICEON
the spe	ecification of which			
(a)[]	is attached hereto	).		
(b)[]	was filed on	a	s Application Serial No.	and was amended
(c) [X]	was described an and amended on	d claimed in Inter	rnational Application No. <u>PCT/US99/</u>	10065 filed on May 7, 1999
includir informa	ng the claims, as a ation which is mate	reviewed and uncomended by any a rial to the patenta	edgment of Duty of Disclosure derstood the content of the above ide mendment referred to above. I acknowled to the subject matter claimed in Regulations § 1.56(a).	owledge the duty to disclose
365(c) insofar States acknow	of any PCT interna as the subject mat or PCT international viedge the duty to den the filing date of	tional application ter of each of the al application in t tisclose material	35 U.S.C. § 120 nited States Code, § 120 of any Unit designating the United States of An claims of this application is not discle he manner provided by the first para- information as defined in 37 CFR § 1 ion and the national or PCT internation	nerica, listed below and, osed in the prior United graph of 35 U.S.C. § 112, I I.56 which became available
(Applicat	ion Serial No.)	(Filing Date)	(Status)(patented,pending,abandoned)	(Patent No. if applicable)
(Applicat	ion Serial No.}	(Filing Date)	(Status)(patented,pending,abandoned)	(Patent No. if applicable)
			Power of Attorney	
Nancy Box 50	J. Parsons, PTO R 68, Alpine Bank Ce	eg. No. 40,364 o enter, 2 <sup>nd</sup> Floor, 2	NO. 32,746, Marina T. Larson, PTO of the firm of OPPEDAHL & LARSON 56 Dillon Ridge Rd., Dillon, CO 8043 of in the Patent and Trademark Office	LLP, having office at P.O. 35 as attorneys to prosecute
SEND	CORRESPONDENCE	7o:	DIRECT TELEPHONE CALLS '	то:
	021121		(970) 468-8600	

PATENT TRACEWORK OFFICE